

List 7

Slope fields, autonomous ODEs, separable ODEs

168. Classify each equation below (the ODE is not shown) as an “implicit solution” or an “explicit solution”.

(a) $x^2 = \sin(3t)$

(d) $y = xe^x - 5x^3$

(b) $y = e^t + C$



(e) $\frac{-1}{x^5} = t^7$

(c) $y = xe^x - 5y^3$

(f) $\ln(y) = 9x$

169. Match the following ODEs to their slope fields.



(a) $\frac{dy}{dx} = \frac{x}{y}$



(b) $\frac{dy}{dx} = \frac{y}{x}$

(c) $\frac{dy}{dx} = \frac{-x}{y}$

(d) $\frac{dy}{dx} = \frac{-y}{x}$



An ordinary differential equation (ODE) for $y(x)$ is **direct** (or **directly integrable**) if it can be written in the form

$$\frac{dy}{dx} = f(x)$$

for some function f . An **autonomous** ODE for $y(x)$ can be written in the form

$$\frac{dy}{dx} = g(y)$$

for some function g . A **separable** ODE for $y(x)$ can be written in the form

$$\frac{dy}{dx} = g(y) \cdot h(x)$$

for some functions g and h .

170. Classify each ODE as “direct” or “autonomous”, and then solve it.

(a) $y' = x^2$

(b) $y' = y^2$

(c) $y' = t^2$

(d) $x' = x^2$

For part (b), you can assume $y = y(t)$ or you can assume $y = y(x)$. It is not clear from the ODE what the input variable is.

171. “Every directly integrable ODE is separable.” Either use formulas to explain why this is true *or* give an example that shows this claim is false.

172. “Every separable ODE is autonomous.” Either use formulas to explain why this is true *or* give an example that shows this claim is false.

173. Solve the autonomous ODE $x' = e^x$.

☆174. Solve the autonomous ODE $y' = \sin(y)$.

175. Solve the autonomous ODE $y' = k y^2$. Your answer should be an explicit general formula for y , but it will also use the letter k .

176. (a) Solve the autonomous ODE $y' = \frac{1}{y}$.

(b) Solve the autonomous IVP $y' = \frac{1}{y}$, $y(1) = 3$.




(c) Solve the autonomous IVP $y' = \frac{1}{y}$, $y(0) = -1$.

☆(d) Solve the autonomous IVP $y' = \frac{1}{y}$, $y(-1) = 0$.

177. Solve the separable ODE $y' = 2^{y+t}$. This is an ODE, so you should give the general explicit solution, if possible.







178. Solve the separable IVP $(1 + x^3) \cdot y' = x^2 y^2$, $y(0) = -1$. This is an IVP, so you should give the particular explicit solution, if possible.

179. One of the three slope fields below corresponds to a directly integrable ODE. Which one?



180. One of the three slope fields from Task 179 corresponds to an autonomous ODE. Which one?

181. For each of the slope fields below, state whether the associated ODE is directly integrable, autonomous, both, or neither.



182. *Newton's Law of Cooling* states that the rate at which an object cools is proportional to the difference in temperature between the object and its environment. As an equation,

$$\frac{dy}{dt} = k(T_{\text{env}} - y)$$

where k and T_{env} are constants. Find the general solution to this ODE (the answer $y(t) = \dots$ will have t , k , T_{env} , and a new constant C in the formula).

183. A hot drink is cooling down according to Newton's Law of Cooling. With the external temperature at a constant 10°C , the drink has cooled from 90° to 85° in 4 minutes. How long will it take to cool down to 60° ?



184. A boat is moving upstream, so the water applies a force $F(t) = -m x''(t)$ that is proportional to the velocity $v(t) = x'(t)$ of the boat. In formulas,

$$x'' = -k x' \quad \text{or} \quad v' = -k v.$$

The boat started its motion with velocity 1.5 m/s , and after 4 seconds it had velocity 1.00548 m/s .

- (a) What distance had the boat traveled after 4 seconds?
 (b) What is the total distance the boat can go?
185. A cylindrical tank has a hole in the bottom, where the liquid flows out with speed proportional to the square root of the remaining volume of liquid in the tank (that is, $V' = k\sqrt{V}$). At the start the tank was full, and after 5 minutes it is half empty. How long will it take until the tank becomes completely empty?
186. For each ODE below: if it is separable, solve it; if it is not separable, write "not separable".

(a) $y' = \sin(t)\sqrt{y}$

(g) $\cos(y)y' = \sqrt{t \sin(y)}$

(b) $y' = 3t^4y^5$

(h) $\frac{y'}{t^3} = \sin(3t)y^2$

(c) $y' = 4e^{3y} \cos(t)$

(i) $yy' = \sqrt{\sin(t)} \cos(t)e^{-y^2}$

(d) $y' = \frac{y^2 + 1}{yt}$

☆(j) $y' = (y^2 - y - 1)t$

(e) $yty' = 1 + y^2$

☆(k) $y' = \frac{\sin(t)}{\sin(y)}$

(f) $y' = 3 \ln(y)t^2y$